



## About CYPRESS

We are researching the investigational drug (amprelosetine) in the Phase 3 CYPRESS trial to confirm whether the investigational drug is effective and safe in treating nOH symptoms in patients with MSA and to learn about any unwanted side effects.



# CYPRESS

An amprelosetine study for  
nOH in patients with MSA

Theravance  
**Biopharma**   
Medicines That Make a Difference®

## More Information :



[www.cypress-study.com](http://www.cypress-study.com)



+1-855-633-8479



[cypress@theravance.com](mailto:cypress@theravance.com)

You can also learn more by visiting:  
[www.clinicaltrials.gov/study/  
NCT05696717](http://www.clinicaltrials.gov/study/NCT05696717)



# The Phase 3 CYPRESS Clinical Trial

## **NOW ENROLLING!**

## Why Participate?

We're taking another critical step toward the goal of neurogenic orthostatic hypotension (nOH) symptom relief in patients with multiple system atrophy (MSA).

Symptomatic nOH affects approximately 70-90% of those with MSA, with symptoms such as dizziness, light-headedness, and feeling faint interfering with one's ability to perform daily activities — especially those requiring standing or walking.

The Phase 3 trial involves 9 visits over a 20-week study treatment period, followed by an optional 2-year long-term extension in which all participants receive the investigational drug. Participants may be eligible to participate in the study remotely (e.g., from your home) after an initial in-office study visit.

Please contact us directly to explore how we might address your specific situation. We'd love to hear from you!



Have you been diagnosed with **multiple system atrophy (MSA)** and have dizziness symptoms associated with **neurogenic orthostatic hypotension (nOH)**? If so, you may qualify.



Scan the QR code above or visit [www.cypress-study.com](http://www.cypress-study.com) for more information regarding trial participation. You can also contact us directly at **+1-855-633-8479**.

## What to Expect:

### Study Highlights

- Once daily, single tablet
- All participants receive the investigational drug during the 12-week open-label period, followed by an 8-week treatment with either placebo or investigational drug

### Duration of Study

- 20-week study treatment period
- Participants who remain in the study will receive the investigational drug for up to 2 years as part of a long-term extension of the study

### Study Visits

- Simple evaluations (similar to regular checkups with questionnaires)
- Only 9 visits in the treatment period

*\*Ampreloxetine is not currently approved by any health agency for use in treating symptoms of nOH in patients with MSA.*